

wherein: A is carbon or silicon; E is nitrogen or P(O); R¹ and R³ are each independently hydrogen, alkyl, aryl, aralkyl, heteroaryl, heteroaralkyl, heterocyclyl or heterocyclylalkyl; p is 0 or 1; Z is any combination of 1-12 units selected from 1,4-phenylene and methylene units, which units may be combined in any order, with the proviso that if the LPC is of formula (Ia) or (Ib), then Z contains at least two phenylene or methylene units; t is 1; X¹ is OH, SH, NH₂, COR⁵ or COOR⁴ where R⁴ is selected from hydrogen, alkyl, aryl, aralkyl, heteroaryl, heteroaralkyl, heterocyclyl and heterocyclylalkyl, and R⁵ is halide, heteroaryl or pseudohalide; n is 3 or 4; Y¹ is CH₂, NH, S or O; Y² is selected from CH and N; R¹, R³, X¹, Y¹, Y² and Z are unsubstituted or substituted with one or more substituents each independently selected from Q; and Q is halogen, hydroxy, nitrile, nitro, formyl, mercapto, carboxy, alkyl, haloalkyl, polyhaloalkyl, aminoalkyl, diaminoalkyl, alkenyl containing 1 to 2 double bonds, alkynyl containing 1 to 2 triple bonds, cycloalkyl, cycloalkylalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, alkylidene, arylalkylidene, alkylcarbonyl, arylcarbonyl, heteroarylcarbonyl, alkoxycarbonyl, alkoxycarbonylalkyl, aryloxycarbonyl, aryloxycarbonylalkyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl,



U.S.S.N. 09/484,484
KÖSTER *et al.*
Preliminary Amendment

B¹ concl'd
~~arylamincarbonyl, diarylamincarbonyl, arylalkylaminocarbonyl, alkoxy, aryloxy, perfluoroalkoxy, alkenyloxy, alkynyloxy, arylalkoxy, amino, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, arylaminoalkyl, diarylaminoalkyl, alkylamino, dialkylamino, arylamino, diarylamino, alkylarylramino, alkylcarbonylamino, alkoxy carbonylamino, aryloxy carbonylamino, aryloxy carbonylamino, azido, alkylthio, arylthio, perfluoroalkylthio, thiocyno, isothiocyano, alkylsulfinyl, alkylsulfonyl, arylsulfinyl, arylsulfonyl, aminosulfonyl, alkylaminosulfonyl, dialkylaminosulfonyl, arylaminosulfonyl or diarylamino sulfonyl.~~



7. (Amended) The LPC of claim 6, wherein Z is a group with three or more points of attachment: one to A, E, or the cyclic nucleus, and the others to two or more X¹ groups.

Sub 2 C²
9. (Amended) The LPC of claim 6, wherein: A is carbon and E is nitrogen.



10. (Amended) The LPC of claim 6, wherein the LPC has formulae (IIa) or (IIb):



B³
17. (Amended) The LPC of claim 6, wherein the LPC has formulae (IIc) or (IId):



22. (Amended) The LPC of claim 6, wherein the LPC has formulae (Ie) or (If):



B⁵

29. (Amended) The LPC of claim 6 that is coupled to a photocleavable linker.

B⁶

31. (Amended) A liquid phase carrier (LPC), selected from the group consisting of tetrakis{6,9-diaza-13-[5'-O-(4,4'-dimethoxytriphenylmethyl)-2'-deoxythymidine-3'-O-yl]-2-oxa-5,10,13-trioxotridecyl}methane ((DMT-dT)₄-PE-LPC), 1,3,5-tris{2,5-diaza-9-[5'-O-(4,4'-dimethoxytriphenyl-methyl)-2'-deoxythymidine-3'-O-yl]-1,6,9-trioxononyl}-benzene ((DMT-dT)₃-Aryl-LPC), tetrakis[13-(2'-deoxythymidin-3'-O-yl)-6,9-diaza-2-oxa-5,10,13-trioxotridecyl]-methane (dT₄-PE-LPC), 1,3,5-tris[9-(2'-deoxythymidin-3'-O-yl)-2,5-diaza-1,6,9-trioxononyl]-benzene (dT₃-Aryl-LPC), tris-{3-aza-4,7-dioxo-7-[5'-O-(4,4'-

dimethoxytriphenylmethyl)-2'-deoxythymidine-3'-O-yl]-heptyl}-amine ((DMT-dT)₃-Amine-LPC) and tris[3-aza-7-(2'-deoxythymidine-3'-O-yl)-4,7-dioxoheptyl]-amine (dT₃-Amine-LPC).







6
B
concl'd

32. (Amended) The LPC of claim 31 selected from the group consisting of tetrakis{6,9-diaza-13-[5'-O-(4,4'-dimethoxytriphenylmethyl)-2'-deoxythymidine-3'-O-yl]-2-oxa-5,10,13-trioxotridecyl}methane ((DMT-dT)₄-PE-LPC), 1,3,5-tris{2,5-diaza-9-[5'-O-(4,4'-dimethoxytriphenyl-methyl)-2'-deoxythymidine-3'-O-yl]-1,6,9-trioxononyl}-benzene ((DMT-dT)₃-Aryl-LPC), tetrakis[13-(2'-deoxythymidin-3'-O-yl)-6,9-diaza-2-oxa-5,10,13-trioxotridecyl]-methane (dT₄-PE-LPC), 1,3,5-tris[9-(2'-deoxythymidin-3'-O-yl)-2,5-diaza-1,6,9-trioxononyl]-benzene (dT₃-Aryl-LPC), tris-{3-aza-4,7-dioxo-7-[5'-O-(4,4'-dimethoxytriphenylmethyl)-2'-deoxythymidine-3'-O-yl]-heptyl}-amine ((DMT-dT)₃-Amine-LPC) and tris[3-aza-7-(2'-deoxythymidine-3'-O-yl)-4,7-dioxoheptyl]-amine (dT₃-Amine-LPC).

7
B
Subst
C-4

39. (Amended) A method of solution phase biopolymer synthesis, comprising the steps of:

(a) reacting an LPC with a first monomer N¹; wherein the LPC has formulae (I):



wherein: A is carbon or silicon; E is nitrogen or P(O); R¹ and R³ are each independently hydrogen, alkyl, aryl, aralkyl, heteroaryl, heteroaralkyl, heterocyclyl or heterocyclylalkyl; p is 0 or 1; Z is any combination of 0-12 units selected from 1,4-phenylene and methylene, which units may be combined in any order; t is 0 or 1; X¹ is OH, SH, NH₂, COR⁵ or COOR⁴, where R⁴ is selected from hydrogen, alkyl, aryl, aralkyl, heteroaryl, heteroaralkyl, heterocyclyl and heterocyclylalkyl; and R⁵ is halide, heteroaryl or pseudohalide; n is 3 or 4; Y¹ is CH₂, NH, S or O; Y² is selected from CH and N; R¹, R³, X¹, Y¹, Y² and Z are unsubstituted or substituted with one or more substituents each independently selected from Q; and Q is halogen, hydroxy, nitrile, nitro, formyl, mercapto, carboxy, alkyl, haloalkyl, polyhaloalkyl, aminoalkyl, diaminoalkyl, alkenyl containing 1 to 2 double bonds, alkynyl containing 1 to 2 triple bonds, cycloalkyl, cycloalkylalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, alkylidene, arylalkylidene, alkylcarbonyl, arylcarbonyl, heteroarylcarbonyl, alkoxy carbonyl, alkoxy carbonylalkyl, aryloxy carbonyl, aryloxy carbonylalkyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylaminocarbonyl, diarylaminocarbonyl, arylalkylaminocarbonyl, alkoxy, aryloxy, perfluoroalkoxy, alkenyloxy, alkynyloxy,

U.S.S.N. 09/484,484
KÖSTER *et al.*
Preliminary Amendment

arylalkoxy, amino, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, arylaminoalkyl, diarylaminoalkyl, alkylamino, dialkylamino, arylamino, diarylamino, alkylaryl amino, alkylcarbonylamino, alkoxycarbonylamino, arylcarbonylamino, aryloxy carbonylamino, azido, alkylthio, arylthio, perfluoroalkylthio, thiocyno, isothiocyno, alkylsulfinyl, alkylsulfonyl, arylsulfinyl, arylsulfonyl, aminosulfonyl, alkylaminosulfonyl, dialkylaminosulfonyl, arylaminosulfonyl or diarylamino-sulfonyl;

(b) separating and purifying the product of step (a);

(c) reacting the product of step (b) with a second monomer N^2 , a dimer N^2-N^3 or a trimer $N^2-N^3-N^4$; and


(d) repeating steps (b) and (c) to produce an LPC-bound biopolymer having m monomers, where m is 3 to 100, wherein:

$N^1, N^2, N^3 \dots N^m$ are biopolymer monomers; and
the dimers and trimers comprise the monomers.

40. (Amended) The method of claim 39, wherein the LPC is selected from the group consisting of tetrakis{6,9-diaza-13-[5'-O-(4,4'-dimethoxytriphenylmethyl)-2'-deoxythymidine-3'-O-yl]-2-oxa-5,10,13-trioxotridecyl}methane ((DMT-dT)₄-PE-LPC), 1,3,5-tris{2,5-diaza-9-[5'-O-(4,4'-dimethoxytriphenyl-methyl)-2'-deoxythymidine-3'-O-yl]-1,6,9-trioxononyl}-benzene ((DMT-dT)₃-Aryl-LPC), tetrakis[13-(2'-deoxythymidin-3'-O-yl)-6,9-diaza-2-oxa-5,10,13-trioxotridecyl]-methane (dT₄-PE-LPC), 1,3,5-tris[9-(2'-deoxythymidin-3'-O-yl)-2,5-diaza-1,6,9-trioxononyl]-benzene (dT₃-Aryl-LPC), tris-{3-aza-4,7-dioxo-7-[5'-O-(4,4'-dimethoxytriphenylmethyl)-2'-deoxythymidine-3'-O-yl]-heptyl}-amine ((DMT-dT)₃-Amine-LPC) and tris[3-aza-7-(2'-deoxythymidine-3'-O-yl)-4,7-dioxoheptyl]-amine (dT₃-Amine-LPC).

45. (Amended) A liquid phase carrier (LPC) that has formulae:

U.S.S.N. 09/484,484
 KÖSTER *et al.*
 Preliminary Amendment



wherein: A is carbon or silicon; E is nitrogen or P(O); R¹ and R³ are each independently hydrogen, alkyl, aryl, aralkyl, heteroaryl, heteroaralkyl, heterocyclyl or heterocyclylalkyl; Z is any combination of 1-12 units selected from 1,4-phenylene and methylene, which units may be combined in any order, with the proviso that if the LPC is of formula (Ia) or (Ib), then Z contains at least two phenylene or methylene units; t is 0 or 1; X¹ is OH, SH, NH₂, COR⁵ or COOR⁴, where R⁴ is selected from hydrogen, alkyl, aryl, aralkyl, heteroaryl, heteroaralkyl, heterocyclyl and heterocyclylalkyl; and R⁵ is halide, heteroaryl or pseudohalide; Y¹ is CH₂, NH, S or O; Y² is selected from CH and N; R¹, R³, X¹, Y¹, Y² and Z are unsubstituted or substituted with one or more substituents each independently selected from Q; and Q is halogen, hydroxy, nitrile, nitro, formyl, mercapto, carboxy, alkyl, haloalkyl, polyhaloalkyl, aminoalkyl, diaminoalkyl, alkenyl containing 1 to 2 double bonds, alkynyl containing 1 to 2 triple bonds, cycloalkyl, cycloalkylalkyl, aryl, heteroaryl, arylalkyl,

U.S.S.N. 09/484,484
KÖSTER *et al.*
Preliminary Amendment

B8
concl'd

heteroarylalkyl, alkylidene, arylalkylidene, alkylcarbonyl, arylcarbonyl, heteroarylcarbonyl, alkoxycarbonyl, alkoxycarbonylalkyl, aryloxy carbonyl, aryloxy carbonylalkyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylaminocarbonyl, diarylaminocarbonyl, arylalkylaminocarbonyl, alkoxy, aryloxy, perfluoroalkoxy, alkenyloxy, alkynyloxy, arylalkoxy, amino, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, arylaminoalkyl, diarylaminoalkyl, alkylamino, dialkylamino, arylamino, diarylamino, alkylarylamino, alkylcarbonylamino, alkoxycarbonylamino, arylcarbonylamino, aryloxy carbonylamino, azido, alkylthio, arylthio, perfluoroalkylthio, thiocyano, isothiocyano, alkylsulfinyl, alkylsulfonyl, arylsulfinyl, arylsulfonyl, aminosulfonyl, alkylaminosulfonyl, dialkylaminosulfonyl, arylaminosulfonyl or diarylaminosulfonyl; R^{20} is alkylene, alkenylene, alkynylene, arylene or heteroarylene; k is 2 or 3; and j is 0 or 1.


47. (Amended) The method of claim 39, wherein the monomers are nucleotides, nucleosides, natural or unnatural amino acids, protein nucleic acid (PNA) monomers or monosaccharides.

B9
Subst
C6

48. (Amended) A method of solution phase biopolymer synthesis, comprising the steps of:

(a) reacting an LPC with a first monomer N^1 ; wherein the LPC has formulae (I):

U.S.S.N. 09/484,484
 KÖSTER *et al.*
 Preliminary Amendment



wherein: A is carbon or silicon; E is nitrogen or P(O); R¹ and R³ are each independently hydrogen, alkyl, aryl, aralkyl, heteroaryl, heteroaralkyl, heterocyclyl or heterocyclylalkyl; p is 0 or 1; Z is any combination of 0-12 units selected from 1,2-, 1,3- or 1,4-phenylene and alkylene, which units may be combined in any order; t is 0 or 1; X¹ is OH, SH, NH₂, COR⁵ or COOR⁴, where R⁴ is selected from hydrogen, alkyl, aryl, aralkyl, heteroaryl, heteroaralkyl, heterocyclyl and heterocyclylalkyl; and R⁵ is halide, heteroaryl or pseudohalide; n is 3 or 4; Y¹ is CH₂, NH, S or O; Y² is selected from CH and N; R¹, R³, X¹, Y¹, Y² and Z are unsubstituted or substituted with one or more substituents each independently selected from Q; and Q is halogen, hydroxy, nitrile, nitro, formyl, mercapto, carboxy, alkyl, haloalkyl, polyhaloalkyl, aminoalkyl, diaminoalkyl, alkenyl containing 1 to 2 double bonds, alkynyl containing 1 to 2 triple bonds, cycloalkyl, cycloalkylalkyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, alkylidene, arylalkylidene, alkylcarbonyl, arylcarbonyl, heteroarylcarbonyl, alkoxy carbonyl, alkoxy carbonylalkyl, aryloxy carbonyl, aryloxy carbonylalkyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylaminocarbonyl, diarylaminocarbonyl, arylalkylaminocarbonyl, alkoxy, aryloxy, perfluoroalkoxy, alkenyloxy, alkynyloxy,